

# Spin Injection into GaAs Quantum Wells with Light




Jason Holden, Yusuke Hashimoto, Hiro MuneKata, Imaging Science and Engineering Laboratory, Tokyo Institute of Technology, Yokohama, Japan



Over the course of the summer, I stumbled over the little Japanese that I know, and worked with Pump and Probe Spectroscopy. By injecting either right or left handed circularly polarized light into a GaAs Quantum Well we are able to cause electrons to be excited with spin polarization. Employing Pump and Probe Spectroscopy we are then able to see this electron spin polarization. This spin manipulation technique is much faster than manipulation using magnetic fields (a few orders of magnitude faster) so it may have uses in creating faster computers and other technologies. This study program has allowed me to learn a wealth of theoretical, cultural, and research knowledge.

## Theory




- GaAs is sandwiched between two layers of AlGaAs as shown above

- Molecular Beam Epitaxy (MBE) is used to deposit the very thin layers on to a substrate. I assisted with a few of these growth sessions


- The quantum well allows this process to be viewed at room temperature

- The laser will pass through the top layer because its wavelength corresponds to less energy than the band gap of AlGaAs



- When circularly polarized light hits the sample it promotes only one band's electrons to the conduction band (due to conservation of angular momentum)

- These electron's spins are all polarized along the axis of the pump laser pulse propagation



- The fact that many of the electrons with a certain angular momentum are already excited, causes an imbalance in  $\sigma+$  and  $\sigma-$  (the indices of refraction for right and left handed circularly polarized light respectively)

- This makes an incident linearly polarized beam (which is a combination of right and left handed circularly polarized light) rotate its polarization plane angle in what is called Kerr rotation


## Future Research

- The next step is to be able to depolarize, or amplify in the middle of the process. Allowing us to switch polarization on and off

- We are now doing Pump and Probe Spectroscopy, with a linear pump pulse, on (In,Mn)As and (Ga,Mn)As

- We are also looking at the effects of a magnetic field on these processes

- With liquid He we can cool the sample down to 4 degrees above absolute zero



Data From GaMnAs

## Experimental Setup



- I set up experiments and helped reduce noise in the measurements


- Titanium-sapphire laser with wavelengths between 720nm and 870nm

- Time Resolution ~ 100 femtoseconds

- Kerr Rotation Sensitivity ~ 1 millidegree


- Laser is chopped into pulses

- Pump Pulse is circularly polarized to inject spin



Sample in cryostat with magnets on either side

## Results



- This graph shows the Magnitude of this Polarization via Kerr rotation

- The decay of the Kerr rotation is a result of spin relaxation and recombination relaxation

- Notice that you must have circular polarized light to get an effect

## Conclusion

- At the nanoscale, changing the size of the crystal affects its physical properties, and allows us to see the spin polarization

- We are manipulating spin with light instead of magnetic fields

- My research time has opened my eyes to the vast amount of knowledge and work that makes up the foundation of new technology

- And I made some Japanese friends in the process

